

INSTRUCTIONS:

1. The question paper contains 7 questions each of 10 marks and total 70 marks.
2. Attempt any 5 question.
3. The missing data, if any, may be assumed suitably.
4. Before attempting the question paper, be sure that you have got the correct question paper.
5. Tables/Data hand book/Graph paper etc. to be supplied to the candidates in the examination hall.

- Q.1(a) What is Galerkin method and how do we apply it in finite element methods? [5]
 Q.1(b) Obtain an approximate displacement equation for the simply supported beam shown in Figure 1 using the trial solution $y(x)=A \sin \pi x/H$. Compare the deflection at the center with the theoretical value $y=-0.06415M_0H^2/EI$. The governing differential equation is [5]

$$EI \frac{d^2y}{dx^2} - \frac{M_0x}{H} = 0$$

Evaluate A by requiring the residual to vanish at (a) $x=H/2$, and (b) $x=0.577H$




Figure 1

- Q.2(a) Evaluate the following integrals for triangular elements [5]
 a) $\int l_1^2 l_2^2 l_3^2 dA$
 b) $\int l_1^2 l_2^2 l_3^3 dA$
- Q.2(b) Derive the shape function for 4 noded rectangular Element in natural Coordinate system. And prove the following: [5]
 a) $N_i + N_j + N_k + N_m = 1$
 b) $N_k = 1$ at k^{th} node and 0 on rest of the nodes.
- Q.3 Derive the shape function for triangular element in Cartesian coordinate [10]




Figure 2

Q.4 The residual equation is given by

[10]

$$R_s = \frac{D^{(s-1)}Y_{s-1} + [D^{(s-1)} + D^{(s)}]Y_s + D^{(s)}Y_{s+1} - L \frac{(Q_{s-1} + 4Q_s + Q_{s+1})}{6}}{L} = 0$$

to obtain the nodal displacements for the beam shown in figure. The governing differential equation is

$$EI \frac{d^2\varphi}{dx^2} - M(x) = 0$$

And $M(x)$ is given in the Figure 3. Each element is 300 cm long; $EI=2(10^{10})$ N. cm².




Figure 3

Q.5 Evaluate the following integrals related to the force vector for the rectangular element.

[10]

$$\int_A GN_i N_j dA, \text{ and } \int_A QN_i dA$$

Q.6 Calculate the axial force in each member of the structural system shown in the Figure 4. Take $E=20(10^6)$ N/cm² and $\alpha = 11(10^{-6})/^\circ\text{C}$

[10]




Figure 4

Q.7 Derive bar element equations using potential energy approach.

[10]

.....25/04/2022.....